


Management of Respiratory Events in Athletes

Andrew S. Houghton, MD, CAQ
Allina Health Orthopedics



*Allina Health Sports Medicine Conference
August 5th 2022*

Allina Health 

1

Disclosures

- None



2

Goals and Objectives

- **Goals**

- Outline differential diagnoses for dyspnea in an athlete.
- Define exercise induced bronchoconstriction (EIB) and exercise induced laryngeal obstruction (EILO) and differences in presentation and treatment.
- Provide brief overview of traumatic pulmonary injuries.

- **Objectives**

- At the end of this presentation one should be able to define the most common respiratory issues in athletes, differentiate presentation and treatment options for EIB and EILO, and be able to recognize the clinical presentation of traumatic pulmonary injuries.

Allina Health

3

Differential Diagnosis for Dyspnea

- **Exercise Induced**

- Exercise Induced Bronchoconstriction (EIB)
- Exercise Induced Laryngeal Obstruction (EILO)
- Physiologic exercise limitations
- Shortness of breath with exercise due to lung diseases (other than asthma)
- Upper respiratory infections/rhinitis
- Spontaneous pneumothorax

- **Traumatic**

- Traumatic pneumothorax, pulmonary contusion, hemothorax, pneumo-mediastinum,

- **Metabolic and cardiac diseases (sickle cell, iron deficiency, arrhythmia, congenital heart disease, coronary artery disease, pulmonary embolism, etc.)**

Allina Health

4

Exercise Induced Bronchoconstriction (EIB)

- **Defined as transient airway narrowing that occurs as a result of exercise.**
- **Inflammatory mediators released when airway osmolarity increases during hyperventilation → results in airway edema and bronchoconstriction.**
- **EIB with asthma (EIBa) and EIB without asthma (EIBwa).**
- **Prevalence:**
 - 5-20% in general population, up to 90% in asthmatic subjects.
- **Signs and symptoms are non-specific**
 - Shortness of breath, fatigue, and underperformance used interchangeably by athletes.

Allina Health

5

Asthma Relationship

- **Patients with mild asthma (do not require inhaled corticosteroid treatment), the prevalence of EIB is 70% to 80%, maybe higher.**
 - Interestingly, only 50% of those treated with inhaled corticosteroids have bronchospastic response to vigorous exercise.
- **Asthma with EIB has been shown to significant affect adolescent Qol scores.**



Allina Health

6

Endurance Athletes with EIB

- **“Occupational Asthma”**
- **Endurance athletes have increased minute ventilation through mouth**
- **Swimmers and cold weather endurance athletes have even higher prevalence**
 - Cold dry air as a trigger
 - Chlorine exposure

Allina Health

7

Initial Screening


- **Thorough history regarding symptoms:**
 - General health
 - Exercise specific
 - Environment specific
- **Insufficient evidence for any existing screening tool**
- **Baseline spirometry w/ and w/o short-acting bronchodilator**
 - Assess airway reactivity should be performed on all athletes as the first diagnostic test to rule out underlying asthma

Allina Health

8

Diagnosis

- **Trial of bronchodilator (may lead to misdiagnosis)**
- **Decrease in FEV1 by 10% with:**
 - Dry-air exercise challenge
 - Eucapnic voluntary hyperventilation
 - Endorsed by International Olympic Committee
- **Mild FEV1 10-25% drop**
- **Moderate 25-50% drop**
- **Severe >50% drop**



www.youtube.com

Allina Health

9

Non-pharmacologic treatment of EIB

- **Interval warm up**
 - 10-15min of vigorous exercise results in refractory period ~2 hours.
- **Face mask or scarf**
 - humidifies and warms air
- **Dietary modification**
 - Questionable evidence
 - Low salt diet, fish oil supplements, lycopene, vitamin C

Allina Health

10

Pharmacological Treatment of EIB

- **If present, make sure underlying asthma is controlled**
- **Inhaled short-acting β -agonist (SABA) - 15min before exercise**
- **Leukotriene receptor antagonist - if using SABA daily**
- **Inhaled corticosteroid - if using SABA daily**
 - May take 2-4 weeks for improvement
- **Combination of bronchodilator and inhaled corticosteroid**
 - Especially when airway inflammation present
- **Other considerations:**
 - Mast cell stabilizing agent

Allina Health

11

Can I train through it?

- **Exercise load is related to degree of bronchial hyper-reactivity**
- **In athletes that have stopped intensive training, EIB has been shown to improve.**
- **So in theory.....no**
- **But....there may be a fitness component**
 - Improved fitness allows for decreased ventilation rate and therefore decrease respiratory dehydration.

Allina Health

12

Doping considerations

- **Inhaled Albuterol (SABA), Salmeterol and Formoterol (LABAs) are the only β -agonists allowed by WADA.**
 - Written for "as needed" is deemed not appropriate, needs specific action plan.
 - Oral or injected form is not allowed
 - Clenbuterol is banned for its anabolic properties
 - Urine albuterol 1000 ng/ml is a positive drug screen (albuterol abuse well above recommended inhaled dosage)
- **Systemic glucocorticoids are prohibited in competition unless a TUE is obtained**

Allina Health

13

Exercise Induced Laryngeal Obstruction (EILO)

- **Paradoxical closure of vocal cords during inspiration.**
 - Also referred to as vocal cord dysfunction (VCD)
- **EILO more accurate term**
- **Frequently misdiagnosed as EIB**
 - Up to 25% of those referred for EIB




Baylor College of Medicine

Allina Health

14

EILO Presentation and Diagnosis

- Intermittent shortness of breath
- Voice changes, stridor, cough
- Usually harder to get air in
- Usually improves quickly after cessation of activity (as opposed to EIB)
- May be linked to environmental triggers
- Link to stress and anxiety
- Diagnosis: spirometry, laryngoscopy



AllinaHealth

15

EILO Treatment

- **Reassure (usually best initial step)**
- **Diaphragmatic breathing control/laryngeal exercise under guidance of speech therapist**
- **Sideline Tip:**
 - Teach them to breath in through nose and out through pursed lips
 - Or try panting
- **Heliox in ED for persistent/severe symptoms**

AllinaHealth

16

Sideline Pearls


Exercise Induced Bronchospasm

- Expiratory symptoms/wheezing
- May take >10 min to start symptoms
- Symptoms persist after cessation of activity
- Proper warm up and

Exercise Induced Laryngeal Obstruction

- Inspiratory symptoms, noisy breathing
- Fast onset and quick resolution of symptoms.
- Education is key


Allina Health




17

Upper Respiratory Tract Diseases

- Upper Respiratory Tract Infection
- Allergic Rhinitis
- Exercise Induced Rhinitis




Allina Health



18

Upper Respiratory Tract Infection

- Intense exercise has been linked to increased incidence of URTI's.
- May be a genetic component that allows for an increased risk of infection when exposed to high physical stress.
- Treat symptomatically
- Ephedrine and derivatives prohibited by WADA.



Allina Health

19

Return to Play Considerations

Upper Respiratory Tract Symptoms

- Training below 70% of maximum heart rate
- Focusing on skill training
- Advance activity as tolerated

Systemic Symptoms

- Systemic symptoms such as fatigue
- Fever >100.4
- Cardiac rate at rest increased by 10 beat/min
- No activity until regression of symptoms

Allina Health

20

Allergic Rhinitis

- 10-45% of world's population - even more prevalent in athletes.
- ~20-40% with AR have symptoms of asthma.
- ~80-90% with asthma have symptoms of AR.
- Affects sleep, decreases concentration and performance.
- most frequent allergic triggers include inhalant allergens: dust mites, pollens, animal dander and molds.
- DX: skin testing
- TX: Allergen immunotherapy, antihistamine, intranasal steroids, avoid allergen
- Screening: Recommendations are for those with AR to undergo spirometry and bronchial hyper-responsiveness testing

Allina Health

21

Exercise Induced Rhinitis

- Defined by Alves et al., 2010 as a fall in peak nasal inspiratory flow above 20% from baseline during exercise
- Unknown mechanism – likely inflammatory mediated. (?similar to EILO)
- Exacerbating factors: cold air, chlorine
- Tx: antihistamines or local ipratropium in the form of nasal inhalation, inhaled corticosteroids for chronic symptoms.



ADAM

Allina Health

22

Exercise Induced Anaphylaxis

- **Spectrum of exercise induced symptoms**
 - Pruritus, urticaria, angioedema, wheezing, syncope
- **More prevalent in females, 50% have history of atopy**
- **Can be any activity, but usually endurance sports**
 - Food dependent variant: exercise 30min – 120min after certain food ingestion results in anaphylaxis.
- **Mechanism unclear, but mast cells degranulation plays a role.**
 - Medication, food, or exposure may prime the cells and exercise triggers degranulation.

Allina Health

23

Exercise Induced Anaphylaxis

- **Presents as pruritus, warmth, redness → hives → angioedema → systemic anaphylaxis (wheezing, syncope, N/V, abdominal pain)**
- **Exercise testing in controlled environment, potential for false negatives**
- **Treatment:**
 - Identify triggers, avoid exercise 4-6 hours after eating, avoid exercise in extremely hot or cold weather.
 - Limited benefit from antihistamines.
 - Injectable epinephrine – make sure coaches, teammates, medical staff are aware and able to administer.

Allina Health

24

Traumatic Pulmonary Injury

- **May occur as injury from direct force, barotrauma, or rapid deceleration of organs against chest wall.**
- **Injuries may include:**
 - Pneumothorax
 - Hemothorax
 - Pulmonary contusion



Allina Health

25

Traumatic Pulmonary Injury

- **Pulmonary injury from blunt thoracic trauma is relatively uncommon in athletics**
- **Most common injury:**
 - Pulmonary contusion
- **Symptoms:**
 - Dyspnea, Chest pain, Dysphonia, Dysphagia, Cough, Hemoptysis
- **Management:**
 - Stabilize ABCs, decision to transport based on severity and mechanism of injury

Allina Health


26

Table 1 Clinical signs and symptoms of pulmonary injury			
Pneumothorax	Hemothorax	Pneumomediastinum	Pulmonary Contusion
Dyspnea	Dyspnea	Dyspnea	Dyspnea
Pleuritic chest pain	Chest pain	Retrosternal chest pain Neck pain Dysphonia Dysphagia Odynophagia Cough	Chest pain Hemoptysis
Tachypnea	Tachypnea	Subcutaneous emphysema	Tachypnea
Tachycardia	Tachycardia	Hamman crunch	Hypoxia
Anxiety	Anxiety	Fever	Wheezing
Hyperresonance to percussion	Dullness to percussion		Rales
Diminished breath sounds	Diminished breath sounds		
<i>Tension PTX:</i> Hypotension Hypoxia Tracheal deviation Distended neck veins	<i>Massive hemothorax:</i> Hypotension		

27

Return to Play After Pulmonary Contusion

- No specific guidelines
- Uncomplicated cases resolve within 7 days
- Respiratory symptoms must have resolved
- Chest wall pain is usually inhibitory



Allina Health

28

Return to Play After Pneumothorax

- No specific guidelines
- Gradual goal oriented progression after radiographic resolution ~3-4 weeks.
- No air travel until 2 weeks after radiographic resolution. (risk of tension pneumothorax)




Allina Health

29

Diaphragm Spasm

- **“Getting wind knocked out”**
 - blow to the epigastric region
- Celiac “solar” plexus = diaphragm spasm
- Tx:
 - supine with knees bent, slow deep breaths
 - focus on abdominal breathing
 - should improve 10-15min





Allina Health

30

Pulmonary Embolism

- **Thromboembolism (usually from legs) to the pulmonary vasculature.**
- **SOB, chest pain, tachycardia, leg pain/swelling (DVT).**
- **Risk Factors:**
 - Long travel
 - Family history of clotting disorder
 - Smoker
 - Oral contraceptive pills
 - Recent immobilization or surgery
- **Treatment: emergent evaluation, 3-6 months daily anticoagulation.**
 - Avoidance of sport is usually recommended until completion of anticoagulation







31

Take Home Points

- **Understand difference between EIB and EILO to avoid mismanagement.**
- **“Above the neck” infections symptoms are cleared to play as tolerated.**
- **Systemic infections symptoms need to resolve before return to play.**
- **Traumatic pulmonary injury is rare, but need to be able to recognize and have plan to transport if necessary.**
- **No clear return to play guidelines for traumatic pulmonary injuries, rather case by case basis with goal oriented progression.**





32

References

- Parsons P, et al. An official American thoracic society clinical practice guideline: Exercise induced bronchoconstriction. *Am J Respir Crit Care med* 2013;187(9):1016-1027.
- Bonini M, Silvers W. Exercise-induced bronchoconstriction: background, prevalence and sport considerations. *Immunol Allergy Clin N Am* 2018; 205-214.
- Smoliga J-M, Mohseni Z-S, Berwager J-D, *et al.* Common causes of dyspnoea in athletes: a practical approach for effective diagnosis and management. *Breathe* 2016; 12: 1-16
- Hull JH, Ansley L, Robson-Ansley P, Parsons JP. Managing respiratory problems in athletes. *Clinical Medicine* 2012;12(4):351-356.
- Brukner P, Khan K. *Clinical Sports Medicine*. 3rd ed. McGraw Hill, 2007,p. 823Y31.
- NPJ Prim Care Respir Med. 2018; 28: 31.
- Gałązka-Franta A, Jura- Szoltys E, Wojciech S, Radosław G. Upper respiratory tract diseases in athletes in different sports disciplines. *Journal of Human Kinetics*. 2016; 53:99-106.
- Alves A, Martins C, Delgado L, Fonseca J, Moreira A. Exercise-induced rhinitis in competitive swimmers. *Am J Rhinol Allergy*, 2010; 24: 114-117
- Felden J. Closed Lung Trauma. *Clin Sports Med* 2013. 32: 255-265

Allina Health

33

Thank You!

Andrew S. Houghton, MD, CAQ
Primary Care Sports Medicine, Non-Operative Orthopedics

952-946-9777
www.allinahealth.org

Allina Health

34