


## When To Call Your Health Care Provider


Call your health care provider if you have any of the following:




- gain 3 pounds in 1 day or 5 pounds in 1 week



- more short of breath than usual



- more swelling of your feet, ankles, legs or stomach than usual



- feeling more tired than usual or being unable to do your everyday activities

- breathing becomes harder when you lie down (You need to start sleeping in a chair.)
- if you are unable to take your medicine as directed

## When to Call 911 or Go to a Hospital Emergency Department

Call 911 or go to a hospital emergency department if you are:

- struggling to breathe
- having chest pain
- confused or unable to think clearly.




# Heart Failure Action Plan

## Do Your Checklist Every Day:

- Weigh yourself in the morning before breakfast. Write down your weight. Compare it to yesterday's weight.
- Take your medicine as directed.
- Check for swelling in your feet, ankles, legs and stomach.
- Eat foods low in sodium (salt). Limit salt to 1,500 to 2,000 mg.
- Be able to do your regular activities without being short of breath.
- Have enough energy to get through the day.


## Call Your Health Care Provider if You Have Any of the Following:




- gain 3 pounds in 1 day or 5 pounds in 1 week



- more short of breath than usual



- more swelling of your feet, ankles, legs or stomach than usual



- feeling more tired than usual or being unable to do your everyday activities

- breathing becomes harder when you lie down (You need to start sleeping in a chair.)

- if you are unable to take your medicine as directed

Whom to Call

Name:

Number:






## Call 911 or Go to a Hospital Emergency Department if You Are:

- struggling to breathe
- having chest pain
- confused or unable to think clearly



# Managing Your Heart Failure — At a Glance

## For an Emergency, Call 911

Daily Weight	Limit the Sodium (Salt) You Eat	Medicines	Activity/Exercise	Call Your Health Care Provider if:
 <ul style="list-style-type: none"> <li>■ Weigh yourself every morning after you urinate and before you eat or drink anything.</li> <li>■ Use the same scale.</li> <li>■ Weigh yourself without clothes.</li> <li>■ Keep a log of your weights.</li> <li>■ Call your health care provider right away if you have a weight gain of 3 pounds in 1 day or 5 pounds in 1 week.</li> </ul> <p><b>Allina Health</b> allinahealth.org</p>	 <ul style="list-style-type: none"> <li>■ Do not use sodium (salt) at the table or for cooking.</li> <li>■ Limit sodium to 1,500 to 2,000 mg each day.</li> <li>■ Read food labels to find out the milligrams of sodium in one serving. Low sodium means 140 mg or less per serving.</li> <li>■ Avoid foods that are high in salt, such as canned items, processed meats (hot dogs, sausage, ham), pickles, olives, prepackaged or convenience foods, fast foods.</li> <li>■ Ask your health care provider about using salt substitutes.</li> </ul>	 <ul style="list-style-type: none"> <li>■ Take your medicines as prescribed each day.</li> <li>■ Carry a current list of your medicines with you at all times.</li> <li>■ Avoid over-the-counter medicines, supplements, and herbal or natural products unless they are recommended by your health care provider.</li> <li>■ Call your pharmacy at least 1 week in advance to refill prescriptions.</li> </ul>	 <ul style="list-style-type: none"> <li>■ Stay active! Try to exercise every day by walking, biking or swimming.</li> <li>■ Use common sense. Do not exercise outside in hot, cold or windy weather.</li> <li>■ If you become short of breath or develop discomfort, stop and rest. If your symptoms don't go away with rest, call your health care provider right away.</li> <li>■ Do not use tobacco. Resources are available for help with quitting. Ask your health care provider.</li> </ul>	 <ul style="list-style-type: none"> <li>■ you gain 3 pounds in 1 day or 5 pounds in 1 week</li> <li>■ you are more short of breath than usual</li> <li>■ you have more swelling of your feet, ankles, legs or stomach than usual</li> <li>■ you feel more tired than usual or unable to do daily activities</li> <li>■ breathing is harder when lying down</li> <li>■ are unable to take medicines</li> </ul> <div style="border: 2px solid red; padding: 5px;"> <p><b>Call 911 if you:</b></p> <ul style="list-style-type: none"> <li>■ struggle to breathe</li> <li>■ have chest pain</li> <li>■ are confused or can't think clearly</li> </ul> </div>

© 2022 ALLINA HEALTH SYSTEM. TM – A TRADEMARK OF ALLINA HEALTH SYSTEM.  
OTHER TRADEMARKS USED ARE OWNED BY THEIR RESPECTIVE OWNERS.  
THIS BOOKLET DOES NOT REPLACE MEDICAL OR PROFESSIONAL ADVICE.  
IT IS ONLY A GUIDE.  
cvs-ah-43732 (4/15)



# Quick Guide To Managing Heart Failure

You are being treated for heart failure. The information in this quick guide will help you better understand heart failure.

This guide also includes things that you and your health care team can do together to help you feel better.

The goal of your treatment is to make you feel better and make you live longer.

By following your lifestyle and medicine plan, your quality of life should improve. Your heart may even work better.

## Heart Failure

Heart failure means that your heart is not working correctly. The main job of your heart is to pump blood to your body. It is like a pump that pumps water out of a pool. If the pump is weak, not all of the water can be removed. Some of the water stays in the pool.

When your heart is weak, it can't pump the blood as well as it should. Parts of your body can hold extra fluid that isn't being moved very well by your heart.

Fluid build-up in your lungs can cause shortness of breath when you walk, go up stairs or lie down. Fluid build-up in your stomach can cause upset stomach after eating, bloating or swelling.

There are two types of heart failure:

- your heart can't fill up with as much as it should
- your heart can't squeeze out as much blood as it should.

In both cases, your heart isn't pumping enough oxygen-rich blood, and fluid is backing up in your lungs and stomach.

## Causes of Heart Failure

There are a lot of causes of heart failure. Some of the common causes are:

- heart attack
- heart valve problems
- high blood pressure
- kidney failure
- irregular heartbeat
- viral infections (that attack the heart)
- chemotherapy and radiation therapy (to treat cancer)
- pregnancy-related (rare condition that can occur in the third trimester of pregnancy or the first few months after giving birth)
- alcohol use (in large amounts)
- street drug use (such as cocaine or heroin)
- genetic (runs in families)
- obesity (being overweight)
- diabetes.

## Treating Heart Failure

You and your health care team will create a treatment plan together. Treating heart failure is done to:

- keep the amount of fluid in your body at a level that your heart can handle
- start or increase medicines that help your heart pump better.

The goal of treatment is to make you feel better and live longer. Your treatment plan may include:

- taking medicines
- eating healthful foods
- getting regular exercise
- reducing your stress.

By following your treatment plan, it is possible that you can have a good quality of life for many years!



## Maintain a Healthy Weight

Each person has a different amount of fluid his or her heart can handle. In general, most patients have too much fluid and need to lose that fluid weight.

Your health care provider will recommend a healthy weight for you. If you do not lose the fluid:

- you will not feel better
- your medicines won't work as well.

## Find Your 'Dry Weight'

Your "dry" weight is your goal weight. This is how much you weigh when you do not have extra fluid (water) build-up in your body.

## Weigh Yourself Every Morning

To maintain the right amount of fluid in your body, you need to weigh yourself each morning.

- Weigh yourself without clothes.
- Weigh yourself after you go to the bathroom and before you eat or drink anything.
- Use the same scale.
- Write down your weight and symptoms in the Heart Failure Weight Log calendar.

You have fluid weight if you gain:

- 3 pounds in 1 day  
or
- 5 pounds in 1 week.

To lose the extra fluid weight, follow your health care provider's directions.

## Watch How Much You Drink

Do not drink more than 2 liters of liquids each day. This includes water, milk, tea, coffee, soda, ice cream and soup.

**Remember:** you need more fluid coming out each day than what goes in. To maintain your weight, the amount of liquids going in each day has to equal what is coming out.

## Limit How Much Salt You Eat

Limit your sodium (salt) to 1,500 to 2,000 milligrams (mg) each day.

- Sodium causes your body to retain (keep) water.
- If you eat more than 2,000 mg a day, your water pills won't work right or maybe not at all.

**Remember:** low salt does not mean no salt. Your heart needs salt to work right. Without any salt, you also will not feel well.

## Take Your Medicines as Directed

There are five types of medicines that work to either improve heart failure or your quality of life. The goal is for you to be on more than two of them at the highest possible doses. The higher dose you can take, the more likely your heart failure will get better.

- **angiotensin receptor blockers (ARBs):** They relax your blood vessels so your heart pumps easier. You would be prescribed this if you cannot take an ACE inhibitor. Examples are losartan (Cozaar<sup>®</sup>), valsartan (Diovan<sup>®</sup>) and candesartan (Atacand<sup>®</sup>).
- **beta blockers:** They can improve how your heart works. Examples are carvedilol (Coreg<sup>®</sup>), metoprolol (Lopressor<sup>®</sup>, Toprol XL<sup>®</sup>), bisoprolol (Zebeta<sup>®</sup>).
- **angiotensin converting enzyme (ACE) inhibitors:** They allow your heart to pump easier. Examples enalapril (Vasotec<sup>®</sup>), captopril (Capoten<sup>®</sup>), lisinopril (Zestril<sup>®</sup> and Prinivil<sup>®</sup>), quinipril (Accupril<sup>®</sup>) ramipril (Altace<sup>®</sup>). If you can't take ACE inhibitors, you will take an ARB.
- **hydralazine and nitrates:** Together, they relax your blood vessels and decrease how hard your heart has to work to pump blood. Examples are hydralazine (Apresoline<sup>®</sup>), isosorbide dinitrate (Isordil<sup>®</sup>), isosorbide mononitrate (IMDUR<sup>®</sup>) or hydralazine/isosorbide dinitrate combination tablet (Bidil<sup>®</sup>).

You may take these if you are already taking the highest doses of beta blockers and ACE inhibitors, or if you can't take ACE inhibitors or ARBs.

- **aldosterone antagonists:** They block the effects of the stress hormone aldosterone. When your body makes too much aldosterone, it causes your kidneys keep more sodium and water. Examples are spironolactone (Aldactone®), eplerenone (Inspra®).

## Get Regular Exercise

Exercise is good for heart failure. The more you exercise, the better your muscles get at pulling the oxygen out of your blood. This means your heart won't have to work as hard.

Push yourself and do a little more walking each day. Your goal is to walk at least 30 minutes each day.

Cardiac rehab can improve your quality of life. Talk with your health care provider for more information.

## Consider Quitting Tobacco Products

### Did You Know

Smokeless tobacco affects your heart in much the same way as cigarettes. Changing to smokeless tobacco is not a good way to protect your health. The best thing to do for your heart health is to quit all tobacco products.

People who smoke cigarettes are 2 to 4 times more likely to develop heart disease than people who do not smoke. Their risk for heart attack, stroke and circulatory problems also increases.

### How tobacco affects you

Tobacco:

- damages and narrows blood vessels
  - This can lead to infections and risk for stroke.
- increases blood pressure
  - This can lead to stroke or heart disease.
- makes your blood vessels and arteries “sticky,” which can block blood flow
  - This raises your risks for heart disease, kidney disease, retinopathy (eye disease that causes blindness) and nerve damage.
- increases insulin resistance
  - This can raise your blood glucose and lead to diabetes.

## Not Ready to Quit? Consider Taking a Break!

If quitting tobacco seems like too much right now, consider taking a break or a vacation from tobacco use.

This can help you feel better by restoring balance\*.

- Set a goal to stop using tobacco.
- Talk with your doctor for resources or ways to cope with withdrawal symptoms.

If this goes well, maybe you will take more breaks during the year. This could lead to a tobacco-free life!

\*Follow your doctor's directions for medicine, exercise, diet and other activities.

## Benefits of quitting

- Your body responds quickly to quitting:
  - **8 hours:** The carbon monoxide level in your blood drops to normal. The oxygen level in your blood increases to normal.
  - **24 hours:** Your chance of heart attack decreases.
  - **48 hours:** Nerve endings start to grow again.
  - **2 weeks:** Your circulation improves and your lung function increases. (Source: World Health Organization)
- There are many health benefits to quitting. Quitting:
  - lowers your chances of stroke, heart disease and nerve damage
  - may lower your blood pressure.

## Suggestions for quitting tobacco

Studies show that the most successful way to quit uses counseling, medicines and follow-up. Ask your health care provider for more information.

- Prepare to stop.
  - Get support from family and friends.
  - Avoid places where you know you will want to use tobacco.
  - Plan activities to replace using tobacco.
- Choose a day to stop.
  - Get rid of cigarettes, ashtrays and lighters.
- Stop.
  - Stop on the day you planned to stop.
  - Be careful with situations or activities in which you might be tempted to start using tobacco again.
  - Try to keep your focus on today, not the future. Tell yourself, "I am not smoking today."
- Stay stopped.
  - Think positive thoughts. Remember why you decided to stop. Reward yourself.
  - Remember that your craving will pass whether you smoke or not.
  - Do not play games like telling yourself, "One cigarette won't hurt," "I deserve a dip or chew," "I just want to see how a cigarette tastes." Your brain might tell you these things to persuade you to go back to tobacco.



## Product-specific Resources

- financial aid Nicotrol® inhaler
  - 1-844-989-PATH (7284)
  - pfizerrxpathways.com
- Plant Extracts aromatherapy
  - 1-877-999-4236
  - plantextractsinc.com

## Resources for quitting

### Allina Health (if you had a recent hospital stay)

- Tobacco Intervention Program at Abbott Northwestern Hospital
  - 612-863-1648
- Tobacco Intervention Program at Mercy Hospital
  - 763-236-8008
- Tobacco Intervention Program at River Falls Area Hospital
  - 715-307-6075
- Tobacco Intervention Services at Allina Health United Hospital – Hastings Regina Campus
  - 715-307-6075
- \*United Hospital Lung and Sleep Clinic Tobacco Cessation Program
  - 651-726-6200
- \*Penny George™ Institute for Health and Healing (LiveWell Center) tobacco intervention coaching
  - 612-863-5178

### Other

- Quit Partner
  - 1-800-QUIT-NOW (1-800-784-8669) or quitpartnermn.com
  - My Life, My Quit™ (ages 13 to 17): text “Start My Quit” to 36072 or call 1-855-891-9989 to talk with a coach
  - American Indian: 1-833-9AI-QUIT or aiquit.com
  - Spanish: 1-855-DEJELO-YA (1-855-335-3569) or quitpartnermn.com/es
  - asiansmokersquitline.org
- online tobacco cessation support
  - smokefree.gov
- American Lung Association/Tobacco Quit Line
  - 651-227-8014 or 1-800-586-4872
- \*Mayo Clinic Nicotine Dependence Center’s Residential Treatment Program
  - 1-800-344-5984 or 1-507-266-1930

\*There may be a cost to you. Check with your insurance provider.

## What Else You Can Do

There are a few other things you can do to help make your heart healthier:

- Do not drink alcohol.
- Keep all of your clinic and lab appointments.
- Call your health care provider if your weight goes up or if you have any new symptoms.

# Heart Failure

## *Your Guide to Living and Succeeding With Heart Failure*

*Fifth edition*

Developed by Allina Health.

© 2022 Allina Health System

The publisher believes that information in this manual was accurate at the time the manual was published. However, because of the rapidly changing state of scientific and medical knowledge, some of the facts and recommendations in the manual may be out-of-date by the time you read it. Your health care provider is the best source for current information and medical advice in your particular situation.

All rights reserved. No part of this book may be reproduced in any form or by any means, electronic or mechanical, including photocopying, without permission in writing from the publisher.

### Disclaimer

This publication is for general information only and is not intended to provide specific advice or recommendations for any individual. The information it contains cannot be used to diagnose medical conditions or prescribe treatment. The information provided is designed to support, not replace, the relationship that exists between a patient and his/her existing physician.

For specific information about your health condition, please contact your health care provider.

## Your Information

Name: \_\_\_\_\_

Address: \_\_\_\_\_

Phone number: \_\_\_\_\_

Emergency contact and phone number: \_\_\_\_\_

\_\_\_\_\_

Primary doctor and phone number: \_\_\_\_\_

\_\_\_\_\_

Insurance provider: \_\_\_\_\_

Insurance card number: \_\_\_\_\_

**Please bring this book with you to every health care provider appointment.**



# Your Health Care Team

Name	Phone
Health care provider: _____	_____
Specialist: _____	_____
Specialist: _____	_____
Nurse: _____	_____
Other: _____	_____
Pharmacy: _____	_____
Emergency contact: _____	
Allina Health clinic: _____	_____
Eye doctor: _____	_____
Home health nurse: _____	

**\* Note: The term “health care provider” can refer to your family doctor, nurse, nurse practitioner, cardiologist, cardiac rehabilitation specialist, or other health care professional you see on a regular basis to manage your heart failure.**

# Table of Contents

## Introduction

Heart Failure: When to Call for Help .....	3
Heart Failure Action Plan .....	5
Managing Your Heart Failure — At a Glance .....	7
Quick Guide To Managing Heart Failure.....	9
Your Personal Information .....	22
Your Health Care Team .....	23
Introduction .....	26
How To Use This Book .....	26

## Chapter 1: Your Heart and Heart Failure

How Your Heart Works .....	25
Heart Failure .....	26
Types of Heart Failure.....	26
Causes of Heart Failure .....	27
Symptoms of Heart Failure .....	29
Special Tests .....	30
How To Manage Heart Failure .....	31
When To Call Your Health Care Provider .....	32
When To Call 911 .....	32
Managing Your Reaction to Heart Failure .....	33

## Chapter 2: Medicines

What You Need To Know About Medicines .....	37
Medicine Dos and Don'ts .....	37
Bring Your Medicines To Your Clinic Visits .....	38
How To Read Your Prescription Label .....	38
Prescribed Medicines .....	40
Over-the-counter Medicines .....	49
Medicine Interactions .....	50
Keep Track of Your Medicines .....	50

## Chapter 3: Sodium (Salt) and Diet

Good Nutrition Can Help Your Heart Failure .....	53
Why You Need to Control Sodium .....	53
How To Read Food Labels .....	60
Dining Out .....	61

Choosing Frozen Meals .....	62
Why You Need To Control Potassium .....	62
Cookbooks .....	64
Websites.....	65
Smart Phone Apps .....	65
Sample Menu Plans for 1 Week .....	65
Low-sodium Recipes .....	69
Food Comparisons .....	74

#### **Chapter 4: Other Treatments**

Implantable Cardioverter-defibrillator (ICD) .....	79
Biventricular Pacemaker or ICD .....	79
Intravenous (IV) Therapies .....	80
Ultra-filtrations.....	80
Left Ventricular Assist Device (LVAD) .....	80
Defibrillator .....	80
Heart Transplant .....	80
Pulmonary Artery Pressure Monitor .....	81
Palliative Care .....	81
Hospice Care .....	82

#### **Chapter 5: Exercise and Activity**

Benefits of Regular Exercise .....	85
Basic Exercise Guidelines .....	85
Good Exercise Choices .....	86
How to Breathe While You Exercise .....	87
Signs You Should Stop Exercising.....	87
Save Your Energy .....	88
Sexual Activity .....	92
Your Everyday Activities .....	93
Personal Grooming .....	94
Household Chores .....	95

#### **Chapter 6: Tobacco Use**

What's In Tobacco .....	99
What Tobacco Does To Your Heart .....	100
What Secondhand Smoke Does To Your Body .....	101
How Your Body Heals After Quitting Smoking .....	101

Quitting Tobacco Use .....	102
Resources For Quitting .....	103
<b>Chapter 7: Glossary.....</b>	<b>107</b>
<b>Chapter 8: Worksheets .....</b>	<b>113</b>

## Introduction

Heart failure means your heart does not pump blood as well as it should. Although some cases of heart failure cannot be cured, heart failure can be managed through diet, exercise and medicine.

This book will give you information about heart failure and how to manage it. This book is meant to be used as a guide. It does not replace medical or professional advice. It is important to keep every appointment with your health care provider.

If you have any questions about heart failure or your general health, please talk with your health care provider.

## How To Use This Book

Heart failure is a disease you can manage. This book has advice and guidelines to help you live the best life you can — and to help keep you out of the hospital. It is a resource for you and your family.

- Use the worksheets to keep track of your weight, medicines and appointments.
- Learn tips for cooking and eating out while on a low-sodium (salt) diet.
- Learn how to safely exercise.
- Learn about heart failure and treatments.
- Bring the book with you to clinic appointments.

This book can also be overwhelming if you try to read it all at once. Read what’s comfortable for you.

**Remember:** your health care team will work with you to help you feel better. Write down any questions or concerns you have. Be sure to bring your list to your next clinic visit.

Between visits, if something doesn’t feel right, call your clinic. For a list of when to call your clinic or 911, see the “Heart Failure Action Plan” on page 5.