

Allina Health's Child Immunization Schedule

Age	Vaccinations			
2 months	DTaP*-HepB*-IPV*	PCV13*	Hib*	RV1
4 months	DTaP*-HepB*-IPV*	PCV13*	Hib*	RV1
6 months	DTaP*-HepB*-IPV*	PCV13*		
12 months	HepA*	PCV13*		
15 months	MMR*	VAR*	Hib*	
18 months	DTaP*	HepA*		
4 to 6 years	DTaP-IPV*	MMR*	VAR*	
11 to 12 years	Tdap*	MCV4*	HPV9 (2 dose series#)	
16 years	MCV4*			
Recommended every year for everyone 6 months and older	Influenza**	**Children younger than age 9 require two doses of the influenza (flu) vaccine at least one month apart the first time they are vaccinated.		

* Vaccines included in the Minnesota School Immunization Law.

Three doses of HPV9 are needed for teens and young adults ages 15 to 26.

+ One dose of MCV4 is needed for previously unvaccinated first-year college students ages 19 to 21 who are or will be living in a residence hall.

Consider for high risk patients at any age: HepB, HepA, MCV4, PCV13, PPSV23. High risk and immigrant populations: for more information, go to: cdc.gov/vaccines or health.state.mn.us/immunize.

Vaccine Abbreviations

- **DTaP-HepB-IPV:** diphtheria, tetanus and acellular pertussis + hepatitis B + inactivated poliovirus vaccine (Pediarix[®])
- **PCV13:** pneumococcal 13-valent conjugate vaccine (Prevnar 13[™])
- **Hib:** haemophilus influenza type b conjugate vaccine (PedvaxHIB[®])
- **RV1:** rotavirus vaccine, oral (Rotarix[®])
- **HepA:** hepatitis A vaccine (Havrix[®])
- **DTaP:** diphtheria, tetanus and acellular pertussis vaccine (Infanrix[®])
- **MMR:** measles, mumps, rubella vaccine (MMR II[™])
- **VAR:** varicella (Varivax[®])
- **Tdap:** tetanus + diphtheria toxoids and acellular pertussis vaccine, adolescent/adult (Boostrix[®])
- **MCV4:** meningococcal conjugate vaccine, quadrivalent (Menveo[®])
- **DTaP-IPV:** diphtheria, tetanus and acellular pertussis + inactivated poliovirus (Kinrix[®])
- **HPV9:** human papillomavirus vaccine (Gardasil[®]9)
- **Influenza:** quadrivalent inactivated influenza vaccine – IIV4 (FluLaval[™])