

Topotecan (Hycamtin®)

Topotecan is an anti-cancer medicine that stops cell growth and division. This medicine is used to treat ovarian cancer and other types of cancer.

How Is It Given?

- clear liquid by IV (intravenous)

What Side Effects May Occur?

This is a partial list of the side effects you may have with this medicine. You may have some, all or none of those listed, or you may have other side effects not listed. Most side effects last only a short time.

Common:

- lowered red blood cell, white blood cell and platelet counts (10 to 14 days after treatment)
- nausea and vomiting
- hair thinning
- loss of appetite, weight loss
- diarrhea
- constipation
- abdominal pain
- headache
- fatigue

Less common:

- muscle or joint pain
- hair loss
- dry mouth
- yellowing of your skin or eyes
- diarrhea — may occur shortly after receiving the drug and can last two to three days
- skin rash, itching
- dizziness

Special Notes

- Topotecan is usually given five days in a row every four weeks.
- You may have changes in your menstrual cycle or sperm count. Discuss birth control measures with your doctor. You should not get pregnant or father a child while taking this medicine.

When To Call Your Doctor

You should call your doctor if you have:

- fever, chills or sore throat
- uncontrolled nausea, vomiting or diarrhea
- unusual bleeding or bruising
- yellowing of your skin or eyes.

For more complete drug information or for questions about this medicine, please ask your doctor, nurse or pharmacist.



ALLINA
Hospitals & Clinics

www.allina.com

© 2009 ALLINA HEALTH SYSTEM
ALLINA® AND THE ALLINA LOGO ARE REGISTERED TRADEMARKS OF ALLINA HEALTH SYSTEM
OTHER TRADEMARKS USED ARE OWNED BY THEIR RESPECTIVE OWNERS

can-ahc-10725 (5/00)